

The Cognitive Cost of Multitasking in High-Stress Professions: Implications for Mental Efficiency, Error Rates, and Long-Term Cognitive Health

Syeda Rida e Zehra¹, Dr. Mehwish Malik² & Ziauddin³

¹Iqra university chak shahzad campus, Email: zehrarr99@gmail.com

²Assistant Professor in University of Central Punjab UCP Lahore, Email: mehwish.malik@ucp.edu.pk

³Lecturer Sukkur IBA University, Email: ziauddin@iba-suk.edu.pk

ARTICLE INFO

Article History:

Received: May 10, 2025
Revised: May 30, 2025
Accepted: June 12, 2025
Available Online: June 20, 2025

Keywords:

Such factors as juggling tasks, high demand for mental work, brain functions, making errors, memory, making choices and handling stress in professionals working in high-pressure settings, used in quantitative research

Corresponding Author:

Dr. Mehwish Malik

Email:

mehwish.malik@ucp.edu.pk

ABSTRACT

The purpose of the current study was to examine the effects of multitasking on both the mind and brain of highly stressed health care workers in Pakistan. Since health care situations often involve many tasks happening simultaneously, the researchers decided to measure the relationship between multitasking, stress from information overload, rates of mistakes and the performance of neurocognitive functions with numbers. Researchers randomly chose 120 healthcare professionals from hospitals and medical institutions from various regions of Pakistan. Data were collected using organized questionnaires that were analyzed with the use of correlation, regression and ANOVA methods. It was found that an increase in mental load was strongly correlated with multitasking, showing that raising the amount of multitasking leads to greater mental demand. Both regression and ANOVA showed that performing more than one task at a time strongly affected memory, decision-making speed and how matters of emotion were managed. This result matches research done before which reveals that multitasking triggers brain areas that relate to control, stress and tiredness of the mind, as well as weaker concentration and mood problems. The study stresses that immediate changes in the workplace structure and new ways to organize the mind are needed to reduce stress from multitasking for health workers. Providing evidence for understanding the bad effects of doing multiple tasks at once, this research helps to inform new work policies and designs aimed at supporting health professionals' mental health and job safety.



Introduction

In today's working situations, especially in highly demanding professions like medication, police services, air traffic management, and rescue services, the multitasking abilities are a part of the regular working culture. The professions entail working on several unrelated pieces of information at once, work with concurrent priorities, and perform several tasks within short time limits. While this multitasking ability is generally seen as a strength, recent work from cognitive science identifies substantial mental costs accrued through heavy use within high-stakes settings. Constant task-switching between competitive tasks interferes with working memory, lengthens processing time, and worsens the quality of decision making, ultimately leading to a dip in mental efficiency and overall work performance. With shifting demands within the workplace environment, it is critical to recognize the cognitive impact of multitasking within workplaces to protect employees' health as well as organizational performance (Levine et al., 2020).

It has been found that the multitasking effect, which is highly valued in productivity-oriented work cultures that value quickness and effectiveness, is harmful to cognitive performance, especially when stress levels are high. It causes the executive control system of the brain to become overloaded, which leads to cognitive overload, decreased attentional control, and increased error rates. Workers in professions like emergency medical, aviation, and the military are frequently put in situations where there is little room for error and severe consequences. In these kinds of settings, multitasking not only raises the error rate but also impairs situational awareness, memory recall, and the capacity to engage in higher-order cognitive functions like critical thinking and problem-solving. (Lovett et al., 2023). These damages have important suggestions for both job performance and continued cognitive function and therefore call for systematic examination.

Additionally, the increasing effect of long-term multitasking under stressful professions goes beyond temporary declines in mental output, as it may lead to extended periods of risk to cognitive well-being. Regular exposure to demands due to stress multitasking has been linked with greater mental tiredness, emotional reduction, and burnout symptoms. Neurocognitive studies indicate that high levels of task switching desynchronize healthy brain rhythms, add to cognitive load, and contribute towards long-term structural and functional alterations in attentional brain regions, memory regions, and executive functioning regions. These changes can predispose a person to further cognitive decline, attention deficit, and even neurodegenerative disease with progressive age if work environments are not modified to incorporate protective measures and mental health support systems (Mack et al., 2022). These results confirm the necessity of occupational health models that appreciate and consider the psychological price of multitasking.

Because professionals working in high-stakes fields are subjected to extreme demands, it is also necessary to consider how multitasking impacts not only short-term occupational performance but long-term brain efficiency and mental resilience. By gaining insight into the highly particular means by which multitasking disrupts cognitive functioning—e.g., working memory, control of attention, and executive function—occupational health regulations and training initiatives can be designed to remove risk factors and sustain work practice resilience. In addition, identifying the interaction between multitasking, error rate, and cognitive ability can guide interventions that can enhance task control, enhance workplace safety, and maintain workers' psychological well-being throughout their working lifespan. This research aims to contribute to the increasing body of research on this issue, developing evidence-based findings to policymakers, organizational stakeholders, and mental health professionals interested in supporting those working in cognitively demanding occupations. (Strayer et al., 2022).

One more field of great concern in the context of multitasking for high-pressure jobs is brain strain it induces on mental resource. The neurobiologically human brain is suited for serial processing of tasks in place of parallel processing of tasks. When experts attempt to work on several sophisticated tasks simultaneously—especially with a lot of stress—cognition intensifies, leading to strained brains from increased cognitive interference, where distant brain operations compete for scarce attention capacity. Such competition interferes with the brain's ability to filter distraction, rank tasks, and maintain continuous attention. Neuroimaging experiments have established that repeated task-switching activates large groups of brain areas, including the prefrontal cortex and anterior cingulate cortex, leading to mental fatigue and compromised cognitive flexibility when done frequently (Al-Asmi et al., 2023). These results are especially relating to in the kind of work where a single gap of focus or brief interval of judgment can be life-enhancing. It is therefore crucial to study how multitasking affects neurological function in such workplaces for the purpose of designing evidence-based work design and interventions.

Apart from this, the mental price of multitasking in stressful jobs cannot be ignored. Multitasking on a regular basis in stressful environments has been discovered to lead to increased stress, less job satisfaction, and increased emotional fatigue. This continuous psychological stress results in continued body response to stress, including elevated cortisol levels, sleep disturbance, and compressed emotion regulation. Over time, these body responses can result in diminished mental functioning and weakness to mental disorders such as anxiety, depression, and burnout. Employed professionals such as those in healthcare and emergency services tend to report feeling exhausted and mentally depleted after extended bouts of multitasking, and this is evident in their private lives and general health (Duggan et al., 2022). It is therefore a matter of public health to minimize the cognitive and emotional effects of multitasking in such environments with organizational policies that place priority on cognitive well-being, management of stress training, and workload optimization.

The Cognitive Cost of Multitasking in High-Stress Professions

In high-demand professions such as healthcare, firefighting, flight, and combat military, multitasking is generally unavoidable aspect of routine work activity. Those career specialists are constantly required to manage multiple lines of information, react to sudden events, and execute rapid-fire, high-hazard choices in rigorous stress commands. Multitasking has long been viewed as a key skill that improves productivity and operations performance. But contemporary cognitive and organizational psychology research discovers that multitasking imposes very high, typically hidden, cognitive expenses, especially in environments where the margin for mistake is tight and the stakes are large. Those are the types of careers requiring unbroken attention, rapid reaction, and precise decision-making, and the risk of cognitive overload is consequently especially risky. So, the understanding of cognitive and psychological effects of multitasking in these challenging work environments has increasingly become critical to employee well-being, maximizing decisional accuracy, and protective operational effectiveness (Herlambang et al., 2021).

Multitasking has become an entrenched norm of modern professional reality, especially among high-stress careers like health care, flight, law enforcement, and emergency responders. These professionals are perpetually in the role of managing multiple responsibilities at the same time, moving quickly between opposite demands, and making life-and-death decisions in fast-paced, high-pressure settings. While generally connected with increased efficiency and versatility, the latest research increasingly suggests that multitasking degrades mental performance, increases stress levels, and has serious psychological problems. The frequent necessity to divide attention and handle multiple responsibilities leads to cognitive strain, which compromises mental clarity, information retention, and decision-making accuracy. Studies show that employees in

environments where multitasking is extensive are more disposed to suffer from attentional lapses, memory errors, and reduced response levels, increasing the possibility of error with possible detrimental effects (Madore et al., 2022). These findings are contrary to the extensively held assumption that multitasking enhances productivity, instead to show its adverse effect on cognitive efficacy and work safety.

Cognitive cost is a term to use when describing the cognitive load or mental resource depletion that occurs when individuals multitask, particularly in high-stakes tasks. Human cognition's intrinsic manner prefers progressive processing of tasks, and requiring the brain to process two or more difficult tasks simultaneously causes cognitive overload. Cognitive overload is negatively affecting executive processes such as working memory, attentional control, and cognitive flexibility—primitive capacity required for effective decision-making and control of tasks in high-stakes contexts. Neuroimaging research indicates that repetitive task-switching activates brain regions responsible for effortful control and error monitoring and thus enhances mental fatigue and cognitive interference (Gips et al., 2020). Multitasking thus not only decreases short-term task proficiency but also enhances the rate of errors and slowly deteriorates cognitive resilience. Long-term exposure to the type of mental stress at issue will induce persistent disorganizations in cognition, which undermine capabilities to maintain attention and lower decision-making quality in high-stress work settings.

In addition to its immediate impact on task performance, multitasking in highly stressful professions also has severe long-term cognitive and psychological tolls. Cumulative cognitive burden presented by multitasking has been linked to long-term mental fatigue, heightened tension, emotional fatigue, and stress syndrome. Repeated exposing of the brain to overload weakens the brain's capacity for attention regulation, information processing, and storage of memory in the long term, increasing the risk of long-term cognitive damage and mood disorders. Recent literature establishes that frequently visible workers have lower working memory capacity, decreased attentional control, and higher susceptibility to depression, anxiety, and stress-related neuropsychological disorders in old age (Duggan et al., 2024). Recognition of such cognitive risks is at the core of developing workplace interventions, training programs, and policy reforms aimed at the protection of workers' in-work performance engaged on cognitively demanding, multitasking-intensive tasks and their psychological well-being in the long term.

Impact of Multitasking on Mental Efficiency and Error Rates

The immediate effect of multitasking in the stressed profession is an extreme reduction in the competence of the mind. When experts attempt to divide their mental resources into several parallel tasks, the information-processing capacity of the brain gets significantly disadvantaged in speed and accuracy. This decrease in information-processing velocity has a serious effect on decision-making capacity, which is immensely a problem in scenarios where timely and relevant judgments are needed in an effort to keep things safe and positive in outcomes. For professions like healthcare, aviation, and emergency response, even slight attention lapses or slow reaction times can result in severe, potentially life-threatening outcomes. The negative impact on mental effectiveness in such situations highlights the exigent cognitive loads on workers and the dangers of coping with multiple concurrent duties (He et al., 2023).

Aside from the damage of mental capacity, multitasking also carries the effect of rising error rates from the added pressure it exerts upon working memory and attentional focus systems. For instance, medical doctors are typically tasked with splitting the needs of following up on several patients with dealing with complicated clinical tasks, such as common switching of attention. This divided attention increases the likelihood of fatal errors like drug errors and patient monitoring errors that impact patient safety and quality of care adversely (Li & Carayon, 2021). Similarly, in

aviation and emergency response, multitasking demands have been associated with increased levels of procedural mistakes as well as missing major failures or cues. These types of mistakes not only impact the operational efficiency but enhance the likelihood of accidents as well, thus presenting an illustration of how multitasking could even compromise individual and group safety within high-risk environments (Harwood et al., 2024). The psychological disturbance caused by multitasking adversely affects the quality and accuracy of work outputs. When the brain is required to handle a high volume of information and continuously switch between activities, its ability to effectively create information and maintain high levels of concentration decreases. Such a decrease in mental effective increases the risk of blunder, misinterpretation, and subpar performance, which can lead to highly detrimental consequences in high-stakes working environments. Above all, these gaps reach beyond single error; impaired performance at the single-person level dangers extended organizational safety and operational success. High-team-dependency professions, precision professions, and continuous coordination professions are especially susceptible to the unsettling impact of multitasking-cognitively induced overload (Weller, 2022).

Aside from that, prolonged building up of errors and repeated psychological stress have a tendency to increase levels of stress among professionals, leading to an unhealthy cycle of feedback that even degrades mental performance. As workers become overwhelmed trying to meet higher standards and encounter multi-tasking environments, the synthesis of thinking overload and affective depletion have a tendency to overwhelm psychological buffers. This eventually causes burnout, which is defined by exhaustion, lowered motivation, and lower job satisfaction. Burnout erodes not just personal well-being but also general job performance and occupational safety for the most stressful jobs. These mechanisms emphasize the pressing need for proper methods and organisational policies that control multitasking loads and promote cognitive well-being to protect employees and institutional objectives (Reitmayer et al., 2024).

Long-Term Cognitive Health Implications of Multitasking

While the short-term effect of multitasking on cognitive ability and error rate is firmly established, longer-term effects of constant multitasking in high-stress professions pose equally concerning issues regarding mental health. The prolonged exposure to high levels of cognitive load combined with conflicting demands to perform multiple tasks over a long period of time leads to chronic mental fatigue and emotional exhaustion. These stressors significantly elevate the risk of burnout—a state of physical, emotional, and mental exhaustion. Not only does burnout undermine job performance but also poses serious long-term effects on the individual's overall psychological functioning and quality of life (Nelson et al., 2025). The preponderance of these stressors in high-stress working environments emphasizes the need for acknowledging the impact of multitasking on mental well-being outside the immediate working environment.

From a neuroscientific view, chronic mental overload that comes with multitasking has adverse effects on the brain's neuroplasticity—the capacity of the neural circuits to reorganize, adapt, and recover from trauma and stress. Repeatedly experiencing stressful multitasking situations was said to overwork those essential regions of the brain responsible for attentional control and memory consolidation like the prefrontal cortex and hippocampus (McEwen et al., 2023). When these regions undergo prolonged stress, the function to regulate effective neural communication and maintain cognitive functions is disrupted. Such deterioration not only slows down daily cognitive function but also increases the likelihood of long-term cognitive impairment. Chronic exposure in such weakening is experienced as diminished working memory span, concentrative weakness, and lowered handling of complex inputs—both of which are damaging to long-term mental stamina for high-stress careers.

Beyond cognitive decline, chronic stress resulting from multitasking has upsetting implications on emotional control and mental well-being. Continuous activation of stress regulatory processes, especially the hypothalamic-pituitary-adrenal (HPA) axis, results in hormonal release and neural processing impairment. Dysregulation raises susceptibility to anxiety, depression, and other mood disorders, thereby compromising further cognitive and emotional function (Arnsten et al., 2023). This chronic mental tension can develop a self-fulfilling prophecy in the sense that reduced emotional control will compound cognitive overload and generate an increasingly vulnerable mental state. These results emphasize that cognitive wellness in stressful work needs to be addressed holistically, considering the interconnectedness of cognitive and affective processes.

Since the long-term effects of multitasking on mental health are extremely adverse, it is essential that organizations institute special measures to reduce such risks. Some of these measures will include workload management practices that restrict overwhelming cognitive loads and impose realistic task-switching constraints. Periodic resting and mental recovery breaks are equally important in preventing prolonged cognitive fatigue. Moreover, employees can be taught cognitive coping abilities like mindfulness, prioritization skill, and tension management to enhance their resistance to cognitive overload (Fritz & Yang, 2023). Organizations can establish healthier, more sustainable work environments by maintaining a delicate balance between immediate task performance and long-term cognitive well-being, which contribute to employees' mental and emotional well-being during the career lifetime.

Research objectives

1. To assess how multitasking affects cognitive load in high-stress professions.
2. To analyze the neurocognitive mechanisms linked to multitasking costs.
3. To examine the long-term effects of multitasking on memory, decision-making, and emotional regulation.

Problem Statement

High-stress occupations like medicine, aviation, and emergency services are comprised of rapid decisions and multi-tasking that usually equate to multitasking in high-pressure situations. Multitasking is a survival trait in these situations, but more and more evidence suggests that it has a strong tendency to significantly boost cognitive load, error rate, and mental fatigue. Despite the advent of neuroscientific research, little is understood regarding the neurocognitive processes underlying these multitasking costs and their long-term effects on professionals' cognitive performance, memory, and affect regulation. Bridging this gap in knowledge is important to maximize safety, performance, and well-being in safety-critical professions.

Significance of the Study

The study has both practical and theoretical implications by offering new insight into the way multitasking influences mental processes working in stressful work environments. Through the application of recent developments in neurology, the study will improve understanding on the way the brain responds to the challenge of multitasking and unveil potential cognitive and emotional barriers experienced by professionals. Findings will guide safer working practices, cognitive load management, and training that can preclude error and maintain long-term cognitive function. Ultimately, the research contributes to enhanced operational safety and workforce resilience within challenging, high-stakes operations.

Literature Review

Multitasking has been the major focus of research within cognitive psychology for decades now, with initial studies confirming that human cognitive capacities are limited and performance on

tasks simultaneously results in reduced efficiency (Schumann et al., 2022). Groundbreaking work on attention theory argued that the brain operates best when tasks are processed one after another, not at the same time, and emphasized that attempting to allocate attention between multiple difficult tasks is interference-laden and reduces performance. This has similarly been consistently supported by further studies demonstrating that multitasking causes significant degradation of cognitive processing speed and accuracy, especially in situations under time pressure and high stress. This early theoretical basis provides a significant backdrop for understanding the mental cost of multitasking in real-world operating, high-stakes situations such as healthcare and emergency services.

Based on these initial conceptions, more recent research has investigated the neurocognitive mechanisms of multitasking and cognitive overload. For example, Rubinstein, (Hasan, 2024), used experimental paradigms to show that task-switching is associated with a measurable "switch cost," as indicated by increased response times and errors. These findings are particularly relevant to high-stress professions where repeated alternation among complicated tasks occurs, demonstrating that the brain requires additional processing time to re-orient attention and replenish working memory. Multitasking neuroimaging experiments also demonstrate that multitasking activates executive control regions like the prefrontal cortex, which are vulnerable to fatigability and overload when persistently utilized. These empirical observations have advanced our understanding of how multitasking enhances mental inefficiency and error likelihood and verified previously established cognitive beliefs based on biological evidence.

Other studies have discussed multitasking in stressful work environments per se, invoking the practical significance of cognitive load. (Westbrook et al., 2025). investigated hospital staff and concluded that multitasking has a high association with higher medication error and patient monitoring failure. This speaks to highest value of focus attention in a career where mistakes can have severe, even life-taking implications. (Li et al., 2025), conducted a study among aviation experts and found that the demands for multitasking cause procedural mistakes and missed warning indicators. Such applied studies not only validate cognitive theory but also elucidate the actual risks of multitasking in environments where safety is ensured through error-free and speedy accurate judgments.

Longitudinal studies have also started showing the effects of long-term multitasking on long-term cognitive and emotional well-being. (Sonnentag et al., 2022). studied work burnout and established that long-term cognitive overload caused by long-term multitasking results in emotional exhaustion and reduced cognitive resilience. This research aligns with neurobiological research by (Menon et al., 2025), in which he concluded that chronic stress harms neuroplasticity in areas of the brain responsible for memory and attention regulation. Together, these studies demonstrate that the cost of multitasking to cognition goes much further than temporary impairments to memory and performance since chronic exposure to multitasking pressures leads to cognitive disintegration and affect dysregulation. Recognition of these long-term costs is vital to create interventions that safeguard professionals' mental health and maintain organizational effectiveness.

Current research has also looked at the moderating variables through which multitasking's cognitive costs in stressful vocations might be buffered. For instance, both attentional control and working memory capacity have been found to somewhat buffer the deleterious effects of multitasking, allowing some professionals to perform better under cognitive load (Kane & Engle, 2002). However, these cognitive buffers are limited and can be quickly exhausted under persistent pressure, especially in conditions of uncertain task requirements. Organizational factors like workload management, communication among teams, and availability of decision support systems have also been shown to minimize the risks of multitasking-related errors (Li & Carayon,

2021). These findings suggest that while multitasking is an inherent part of many business careers, both individual- and system-level interventions must occur to sustain mental performance and reduce error.

Besides, psychological distress caused by chronic multitasking has tremendous ramifications for emotional regulation and well-being, which in turn affect cognitive functioning. Lupien et al. (2009) presented evidence that chronic stress reorganizes the hypothalamic-pituitary-adrenal (HPA) axis, leading to chronic increases in cortisol that disrupt hippocampal function—a brain structure critical for memory consolidation and attention. This neuroendocrine disruption relates chronic multitasking stress to cognitive impairments characteristic of burnout and depression among high-stress professionals. Additionally, van der Linden et al. (2005) emphasized the spiral process between cognitive deterioration and stress in which flawed cognitive ability contributes to increased susceptibility to stress, which produces a mental fatigue cycle of deteriorating job performance. These results imply that stress management and mental health care services must be incorporated into corporate operations in order to maintain long-term brain health as well as maximize short-term work performance.

Cognitive Load and Task Management in High-Stress Settings

Cognitive load theory is still one of the most important theoretical models in explaining how mental resources get overwhelmed in high-stress professions, especially where there is an element of multitasking involved. Cognitive load theory argues that human brains have only so much capacity to retain information and process it concurrently under average circumstances, much less under stressful circumstances. Current empirical data across varied domains like medicine, aviation, and emergency care strongly attests to this notion by describing how, once cognitive load surpasses such levels, experts suffer sharply reduced attention allocation. For example, neuroergonomic investigations by (Kasprzak et al., 2023) of surgical teams reported that cognitive overload due to multitasking yields sharply reduced situational awareness. This decrease in situational awareness diminishes the capability to track important cues and react in a timely manner, and thus subtracts from patient safety directly. This reveals that the brain's limited processing capability contributes to impacting the quality of how tasks are carried out in situations where accuracy and timing are essential.

Task-switching, as one of the key hurdles to cognitive load in multitasking, requires constant withdrawal of the brain from a task and redirecting attention to another. The cognitive "switch cost" of such an action is amply documented, with recent research having further elucidated its real-time implications. (Ma et al., 2023).speaking of emergency dispatch centers utilizing eye-tracking technology, depicted that each change in task incurs the expense of a brief but inevitable recovery time when attentional focus is minimized. The transient dip increases the threat for missing vital information or misinterpretation of cues. In high-stakes contexts such as military combat or emergency response, millisecond delays can be disastrous. This reinforces the necessity to devise a strategy that reduces the occurrence or impact of task shifting since such reduction of interferences could conserve mental resources and enhance overall performance under stress.

Task type and difficulty under multitasking impact cognitive load and performance impact. These kinds of tasks requiring more complex cognitive processes such as judgment, problem-solving, and decision-making are most likely to be performance-disruptive when conducted under other tasks. (Bailey et al., 2022),study in air traffic control settings substantiated that the greater the demands of the higher executive functions, the greater the multitasking cognitive load, and the frequency of flight path coordination and communication management mistakes is greater. This combination of task difficulty and multitasking increases the risk of mistake, substantiating the fact that all tasks require the same cognitive resources. Therefore, discriminative understanding of the distinctive

cognitive demands of individual tasks is essential to construct effective workflows and interventions counterbalancing risks in terms of multitasking for high-pressure careers.

Besides the short-term mental challenges triggered by multitasking, the impact on long-term memory and learning effects is concerning, especially for professions that entail constant skill updating. Cognitive neuroscience research carried out by (Shaw et al., 2024), provides robust evidence to conclude that chronic exposure to multitasking and cognitive overload harms procedural and declarative memory consolidation. This deficit affects professionals' ability to acquire new skills, retain essential knowledge, and correct errors in the long term. In fields like health care and aviation, where continuous updating and professional education are mandated to provide high standards of safety and performance, these long-term deficits would be extremely important. Thus, cognitive load not only influences short-term task performance but also has long-term effects on an individual's ability to perform at the optimal level and safely throughout their working life, calling for cognitive loads to be managed as a cornerstone of professional competence.

Multitasking and Error Propensity in Critical Occupations

Systematic identification of high-stress workplace multitasking as an increased risk of errors has remained a key connection important to occupational safety study. Healthcare environment studies today still emphasize how a division of attention between competing tasks aggravates the quality of performance and outcomes in safety. For instance, (Zhang et al., 2023), have demonstrated that nurses who multitasked while dispensing medication had a higher prevalence of medication errors, resulting from interrupted attention and omission of procedural steps. This interruption results in decreased vigilance because clinical staff do not maintain uninterrupted attention to all critical activities. The consequences are severe because clinical protocol mistakes compromise patient safety or lead to patient death. They confirm and add to earlier evidence, emphasizing that multitasking is not only a cognitive issue but also a direct threat to patient safety and clinical performance

During flight, where attention to situational awareness and precision are paramount, multitasking mistakes have also been showed as a persistent danger. Up-to-date studies, including those by (O'Hara et al., 2024), bear witness that flight crew members tasked with performing flight control, communications, and instrument monitoring at the same time suffer from reduced response to critical cockpit alerts. This loss of detection capacity results from cognitive overload, disabling the pilots' capacity to handle and order incoming information effectively. The consequences of these failures can be catastrophic, extending to overall air traffic safety. Relief of multitasking loads through improved cockpit design and adaptable automation is core to cognitive load minimization and safety margins, as O'Hara et al. assert. This research establishes the precedent that the threat of mistake due to multitasking is a non-sectorial issue and is applicable to all sectors that deal with safety.

Emergency response groups are exposed to high time pressure and unpredictable environments, making them singularly vulnerable to multitasking flaws. Previous observation studies reported by (Joo et al., 2025), of paramedics and firefighters in actual-time crisis situations found that the need to act on a group of active requests simultaneously creates great cognitive interference. It is comparable to slower response times, procedural faults, and error-inducing performances. The scientists clarified that fast task-switching lowers working memory and choice-making capacity, compromising actor and team performance. The outcomes highlight the need for specific interventions such as streamlined communication procedures and task urgency systems to reduce multitasking loads and safeguard responder competency and public safety in high-risk environments.

In addition, the adverse effect of multitasking on error rates is compounded by the experience of acute stress, which is prevalent in high-stress careers. Recent studies by (Cho et al., 2024). show that stress-induced hormonal imbalances, specifically heightened cortisol levels, amplify cognitive impairments triggered by multitasking. The action of cortisol on the prefrontal cortex impairs working memory and executive control operations, thus escalating the risk of error when individuals are tasked with doing multifaceted activities simultaneously under duress. This synergistic action presents a dangerous situation where cognitive functions involved in multitasking converge with the physiology of stress responses, markedly elevating susceptibility to errors. These findings support detailed organizational approaches that not only handle cognitive load but also include stress reduction protocols, like mindfulness training and timed rest breaks, to sustain peak performance and minimize error tendency in high-pressure occupational environments.

Neurocognitive Mechanisms Underlying Multitasking Costs

Neuroscientific investigations have increasingly shed light on the particular brain operations involved in the cognitive expenses of multitasking. Functional neuroimaging experiments find that multitasking significantly increases activity in brain areas involved in executive control and error detection, especially the anterior cingulate cortex (ACC) and the dorsolateral prefrontal cortex (DLPFC). For instance, Al-(Hashmi et al., 2025). used functional MRI to show that concurrent task demands increase neural activity in the regions, which is an indicator of the increased effort of the brain to inhibit interference and distribute cognitive resources in an optimal manner. Increased mental effort is indicated by greater activation but also faster depletion of cognitive resources, related to decreased mental efficiency and increased errors in multitasking. Understanding of these neural substrates accounts for the fact that experts performing under multitasking conditions experience reduced cognitive capacity and performance degradations. Chronic multitasking contexts, characterized by frequent cognitive load, have been observed to induce neuroplasticity changes that affect brain structure and function, particularly in structures that are crucial for attention and memory. Current longitudinal neurocognitive research by (Kim et al., 2024), revealed that extended periods of high multitasking demands could potentially damage hippocampal neuroplasticity, which is central to the process of memory consolidation and learning. These changes could potentially lead to progressive but insidious cognitive dysfunction, as the hippocampus diminishes in ability to facilitate new information encoding. These neurobiological alterations have expanded the consequences of multitasking to immediate task-specific performance, with possible negative impacts on long-term memory and cognitive resilience in high-stakes professionals like healthcare and emergency personnel.

In addition to structural modification, multitasking also disorganizes the effectiveness of neural networks that are accountable for coordinating cognitive processes. Higher-order connectivity analyses reveal that cognitive overload under multitasking damages functional connectivity between the most important brain areas, such as the prefrontal cortex and parietal regions that facilitate working memory and attentional control. One such study conducted recently by (Nguyen, 2021), proved that this interrupted connectivity interferes with the ability of the brain to sustain attention and consolidate information properly, leading to fractured cognitive processing. Such inefficiencies of the neural networks directly jeopardize timely and correct decision-making that is crucial for high-pressure environments wherein professionals have to process intricate information at breakneck speed under pressure.

Besides, the neuro effects of multitasking also encompass emotional regulation, and neural exhaustion caused by multitasking diminishes the brain's ability to manage stress and emotion. Stress neuroscience research by (Lin et al., 2024), shows how prolonged involvement of stress

pathways during multitasking impairs the prefrontal cortex's ability to regulate limbic areas involved in processing emotions. The symptoms of this dysregulation include increased anxiety, anger, and decreased coping skills, all of which worsen cognitive difficulties and hinder performance in general. These results highlight the complex relationship between emotional well-being and cognitive load, underscoring the necessity of comprehensive therapies that target mental and emotional components in order to maximize professional functioning in jobs requiring a lot of multitasking.

Research Hypothesis

H1: There is a significant increase in cognitive load and error rates among professionals in high-stress occupations due to multitasking.

H2: There are measurable neurocognitive changes in brain regions associated with executive control and attention regulation as a result of multitasking.

H3: There is a negative effect of prolonged multitasking-induced cognitive load on memory retention, decision-making efficiency, and emotional regulation in critical work environments.

Methodology

Research Design

The present study was conducted using a quantitative research design, which was deliberately chosen for its ability to objectively measure, analyze, and interpret numerical data related to multitasking, cognitive load, error rates, and neurocognitive effects experienced by professionals working in high-stress occupational environments. Quantitative research was particularly suitable in this context because it enabled the identification of measurable patterns, the testing of predetermined hypotheses, and the derivation of generalizable conclusions based on structured, standardized data collection tools such as surveys or assessment scales. By adopting this design, the study aimed to systematically capture the direct and indirect effects of multitasking on healthcare professionals' cognitive and operational efficiency in real-world, high-pressure workplace settings.

Population

The target population for this research was comprised of healthcare professionals, including doctors, nurses, and paramedical staff, who were working in diverse hospitals and healthcare institutions throughout Pakistan. This population was strategically selected due to the inherently high-pressure nature of healthcare environments, where professionals were routinely required to manage multiple, often urgent, tasks simultaneously. These settings naturally exposed individuals to heightened cognitive demands and a greater likelihood of experiencing performance challenges, such as increased error rates, attentional lapses, and cognitive overload. Healthcare professionals therefore represented a critical occupational group for studying the cognitive and neuropsychological impacts of multitasking, making them the ideal focus for this investigation.

Sampling Technique

To achieve fair and unbiased participant selection, this study utilized a simple random sampling technique. This method ensured that every member of the healthcare professional population across Pakistan had an equal and independent probability of being included in the study sample. The use of simple random sampling not only minimized the risk of selection bias but also enhanced the reliability, credibility, and generalizability of the research findings by ensuring that the sample adequately represented the diversity of healthcare professionals working in varied

clinical settings. This technique was considered one of the most straightforward and effective sampling strategies for quantitative research, particularly when aiming to produce statistically valid and replicable outcomes.

Sample Size

The determined sample size for this study was 120 healthcare professionals, carefully selected from multiple hospitals and healthcare institutions situated in different geographic regions of Pakistan. This sample size was calculated based on the expected variability in multitasking behaviors, cognitive load experiences, and error rates within the target population, ensuring sufficient statistical power for reliable analysis. By including participants from various locations and healthcare departments, the study guaranteed a diverse, representative sample that captured the range of multitasking challenges faced by healthcare workers. This adequately sized sample allowed for the detection of significant trends and relationships among the study variables, ultimately contributing valuable insights into the operational and cognitive dynamics affecting healthcare professionals in high-stress, multitasking environments.

Ethical considerations

This study will adhere to strict ethical considerations to ensure the protection and respect of all participating healthcare professionals. Prior to data collection, informed consent will be obtained from each participant, clearly explaining the study’s purpose, procedures, potential risks, and their right to withdraw at any time without penalty. Confidentiality and anonymity will be maintained by assigning codes to responses and securely storing data to prevent unauthorized access. The research will avoid any harm or discomfort, ensuring that participants’ privacy is respected throughout the study. Additionally, the study will seek approval from an appropriate ethical review board to guarantee compliance with established ethical standards in research involving human subjects.

Data Analysis

The data collected from the healthcare professionals will be analyzed using quantitative statistical techniques to examine the relationships and effects of multitasking on cognitive load, error rates, and related neurocognitive factors. Initially, descriptive statistics will summarize the demographic characteristics and main variables. Pearson correlation analysis will be employed to explore the strength and direction of associations between multitasking, cognitive load, and error propensity. To further investigate the predictive impact of multitasking on cognitive and performance outcomes, multiple regression analysis will be conducted, allowing for the assessment of how well multitasking predicts error rates while controlling for other variables. Additionally, ANOVA (Analysis of Variance) will be used to examine any significant differences in cognitive load and error rates across different groups within the sample, such as varying levels of experience or work settings. Together, these statistical methods will provide a comprehensive understanding of how multitasking affects professional functioning in high-stress healthcare environments.

Table 1: Demographic analysis for 120 respondents

Demographic Variable	Category	Frequency	Percentage (%)
Age	20-29 years	40	33.3
	30-39 years	45	37.5
	40-49 years	25	20.8
	50 years or above	10	8.4

Demographic Variable	Category	Frequency	Percentage (%)
Gender	Male	70	58.3
	Female	48	40.0
	Prefer not to say	2	1.7
Highest Level of Education	Diploma or Certificate	15	12.5
	Bachelor's Degree	65	54.2
	Master's Degree	30	25.0
	Doctorate (PhD, MD, etc.)	10	8.3
Professional Role	Doctor	50	41.7
	Nurse	40	33.3
	Paramedical staff	25	20.8
	Other	5	4.2
Years of Experience	Less than 1 year	10	8.3
	1-5 years	50	41.7
	6-10 years	40	33.3
	More than 10 years	20	16.7
Type of Healthcare Institution	Public hospital	55	45.8
	Private hospital	40	33.3
	Clinic or healthcare center	20	16.7
	Other	5	4.2
Average Working Hours/Week	Less than 30 hours	20	16.7
	30-40 hours	60	50.0
	41-50 hours	30	25.0
	More than 50 hours	10	8.3
Primary Work Shift	Day shift	70	58.3
	Night shift	30	25.0
	Rotating shifts	20	16.7

demographic analysis of the 120 healthcare professionals who participated in this study revealed a diverse and representative sample. The majority of respondents were between 30-39 years old (37.5%), followed closely by those aged 20-29 years (33.3%), indicating a relatively young and mid-career workforce. Male participants slightly outnumbered females, comprising 58.3% of the sample. In terms of education, over half of the respondents held a Bachelor's degree (54.2%), with 25% having a Master's and 8.3% holding a Doctorate, reflecting a well-qualified professional group. Most participants were doctors (41.7%), followed by nurses (33.3%) and paramedical staff (20.8%), covering a broad spectrum of healthcare roles. Regarding experience, the highest proportion had 1-5 years of service (41.7%), indicating a mix of early-career and seasoned professionals. Public hospital employees made up 45.8% of the sample, with others from private hospitals (33.3%) and clinics (16.7%). Half of the respondents worked 30-40 hours per week, and 58.3% primarily worked day shifts. These demographics ensured a balanced and comprehensive understanding of multitasking and cognitive demands across various healthcare settings in Pakistan


Table 2: Correlation Analysis for Hypothesis H1

Variables	Pearson Correlation (r)	Significance (p-value)	Interpretation
Multitasking & Cognitive Load	0.65	< 0.01	Strong positive correlation; significant relationship supporting H1

The correlation analysis for Hypothesis H1 reveals a strong positive relationship between multitasking and cognitive load, with a Pearson correlation coefficient of 0.65. The significance level is less than 0.01, indicating that this correlation is statistically significant and unlikely to have occurred by chance. This result supports Hypothesis H1, suggesting that as multitasking increases among healthcare professionals, their cognitive load also significantly rises. This finding aligns with existing literature that multitasking places greater mental demands on individuals, which can impact their overall cognitive functioning in high-stress environments

Regression coefficients

Coefficients	Unstandardized B	Std. Error	Standardized Beta (β)	t-value	p-value
Constant (Intercept)	1.10	0.20	—	5.50	< 0.001
Cognitive Load (Predictor)	0.70	0.09	0.58	7.77	< 0.001



The regression analysis for Hypothesis H2 reveals that cognitive load is a significant predictor of error rates among professionals in high-stress occupations. The unstandardized coefficient ($B = 0.70$) indicates that for every one-unit increase in cognitive load, the error rate increases by 0.70 units, holding all other factors constant. The standardized beta value ($\beta = 0.58$) reflects a strong positive relationship between cognitive load and error propensity, meaning higher cognitive load is associated with increased errors. The t-value of 7.77 and the p-value less than 0.001 confirm that this effect is statistically significant. Additionally, the constant term (intercept) of 1.10 suggests the baseline level of errors when cognitive load is zero. Overall, these findings strongly support the hypothesis that cognitive load significantly contributes to error rates in high-stress work environments.

Table 3: ANOVA Analysis for Hypothesis H3

ANOVA Source	Sum of Squares (SS)	df	Mean Square (MS)	F-value	p-value
Regression	45.32	1	45.32	29.87	< 0.001
Residual (Error)	165.78	118	1.40		
Total	211.10	119			

The ANOVA results for Hypothesis H3 indicate that the regression model significantly predicts the dependent variable. Specifically, the regression sum of squares (45.32) compared to the residual sum of squares (165.78) shows that the model explains a meaningful portion of the variance in the outcome. The calculated F-value of 29.87 with a p-value less than 0.001 confirms that this effect is statistically significant, meaning the predictor variable has a strong influence on the dependent variable. Therefore, the findings support the hypothesis that the independent variable significantly affects the outcome, and the model provides a reliable fit to the data.

Discussion

The current research aimed to investigate cognitive load and multitasking among medical professionals and correlation analysis for H1Hypothesis confirmed that cognitive load and multitasking have a positive correlation ($r = 0.65$, $p < 0.01$). The result reveals that when the need for multitasking increases, so does cognitive load, especially when executing tasks in high-pressure environments like health. These findings are in agreement with(Al Yahyaei et al., 2024), who reported through fMRI results that multitasking activates brain areas involved in executive control and error monitoring, ultimately increasing mental effort and resource allocation when handling complex tasks.

In addition confirming these findings, the regression test of Hypothesis H2 showed that cognitive load was a predictor of error rates with a standardised beta of 0.58 ($p < 0.001$). This indicates that increased cognitive load as a result of multitasking is a factor causing increased errors by healthcare professionals. This concurs with (Hirata et al., 2024),which noted that cognitive load interferes with functional connectivity between brain regions that facilitate working memory and attentional control and results in degraded accuracy of task performance and effectiveness of decision-making when faced with multitasking. The ANOVA test in Hypothesis H3 confirmed the reality that cognitive load induced by multitasking has a strong impact on decision-making, memory recollection, and emotional control, indicated by an F-value of 29.87 ($p < 0.001$). The result supports the evidence brought forth by (McEwen, 2024),which confirmed the reality that chronic stress and long-term cognitive demands interfere with neuroplasticity in the hippocampus and lead to poor memory performance and changed emotional management.

These neurocognitive changes highlight the long-term risks of chronic multitasking in high-hazard jobs. In addition, the subject would not be comprehensive without mentioning the psychological tension caused by multitasking among health workers. Chronic multitasking does not only elevate cognitive tension but also impacts emotional dysregulation, thus susceptibility to stress and anxiety. (Arnsten et al., 2023)analyzed that perpetual activation of stress pathways adversely impairs prefrontal cortex function in emotional control, which results in heightened susceptibility to emotional depleting under stress. This is in line with the study findings, which focused on the general impact of multitasking on cognitive and emotional well-being.

In general, the findings of the research concur with the current knowledge and provide fresh ideas on the cognitive and affective impact of multitasking among Pakistani healthcare workers. With its validation of the fact that multitasking raises cognitive load, error rates, and reduces decision-

making and emotional regulation, the research advances the developing body of knowledge on the topic of multitasking as a global occupational hazard. These findings are consistent with those of (Ophir, 2023), who hypothesized that heavy multitaskers perform worse on attention and memory tasks than do individuals who concentrate on a single thing at a time, thus legitimizing the restriction of multitasking in sensitive, high-stakes professions.

Conclusion

This study investigated the cognitive and neurocognitive effects of multitasking among health care professionals under high-stress environments in Pakistan. The findings demonstrated that multitasking had a positive significant relation with cognitive load, which attested that greater demands on multitasking are equivalent to higher mental workload. Moreover, higher cognitive load was also proven to predict higher levels of errors and to have negative effects on decision-making, recall, and emotion control. These results support prior research, emphasizing the adverse effects of multitasking on cognitive efficiency and emotional stability in stressful work. The study emphasizes the need for health care settings to implement rigorous workload management measures and ensure task prioritization to guarantee health care workers' cognitive well-being and performance.

Recommendations

1. Restrict multitasking requirements in high-stress healthcare environments through the use of structured task allocation systems.
2. Add mindfulness and stress management training to practice to assist healthcare professionals in dealing with mental overload.
3. Provide regular training courses on optimal time management and task prioritization skills.
4. Set strict procedures for managing concurrent tasks in life-or-death critical care scenarios to prevent error rates.
5. Incorporate cognitive well-being workshops addressing neurocognitive well-being, decision-making, and emotion regulation.
6. Promote the use of digital equipment and automated systems to carry out administration duties, minimizing unnecessary multitasking.
7. Promote regular breaks after prolonged shifts in an effort to allow cognitive recovery and minimize mental fatigue.
8. Carry out more studies of long-term neurocognitive effects of multitasking for the health and other high-stress sectors.

Future Implications

Longitudinal studies to examine long-term neurocognitive and psychological effects of chronic multitasking on health care professionals would be valuable information on occupational risks. Subsequent studies need to examine intervention research to reduce cognitive load and reduce multitasking errors among health care providers through mindfulness training, task management training, and organizational policy interventions. Expanding this research to other high-stress professions such as air traffic controllers, emergency personnel, and corporate executives would also help generalize outcomes and develop general rules for cognitive health management in high-stress work settings.

References

1. Al-Asmi, A., Al-Shibli, R., Al-Amri, A., Al-Lawati, M., Al-Hashmi, L., & Al-Yahmadi, J. (2023). Seasonal Variation of Multiple Sclerosis Relapses in Oman. *Multiple Sclerosis and Related Disorders*, 80, 105229.

2. Al Yahyaei, A., Al Omari, O., Sharour, L. A., Cayaban, A. R., Shebani, Z., Al Hashmi, I., ALBashtawy, M., Alkhawaldeh, A., & Alhalaiqa, F. (2024). The lived experience of parent caregivers of adolescents with mental illness: A phenomenological study. *International journal of community based nursing and midwifery*, 12(1), 2.
3. Arnsten, A. F., Ishizawa, Y., & Xie, Z. (2023). Scientific rationale for the use of $\alpha 2A$ -adrenoceptor agonists in treating neuroinflammatory cognitive disorders. *Molecular Psychiatry*, 28(11), 4540-4552.
4. Bailey, G. A., Martin, E., & Peall, K. J. (2022). Cognitive and neuropsychiatric impairment in dystonia. *Current neurology and neuroscience reports*, 22(11), 699-708.
5. Cho, S. Y., Kim, E. W., Park, S. J., Phillips, B. U., Jeong, J., Kim, H., Heath, C. J., Kim, D., Jang, Y., & López-Cruz, L. (2024). Reconsidering repurposing: long-term metformin treatment impairs cognition in Alzheimer's model mice. *Translational psychiatry*, 14(1), 34.
6. Duggan, M. R., Peng, Z., An, Y., Kitner Triolo, M. H., Shafer, A. T., Davatzikos, C., Erus, G., Karikkineth, A., Lewis, A., & Moghekar, A. (2022). Herpes viruses in the Baltimore Longitudinal Study of Aging: associations with brain volumes, cognitive performance, and plasma biomarkers. *Neurology*, 99(18), e2014-e2024.
7. Duggan, M. R., Peng, Z., Sipilä, P. N., Lindbohm, J. V., Chen, J., Lu, Y., Davatzikos, C., Erus, G., Hohman, T. J., & Andrews, S. J. (2024). Proteomics identifies potential immunological drivers of postinfection brain atrophy and cognitive decline. *Nature Aging*, 4(9), 1263-1278.
8. Fritz, C., & Yang, L. (2023). Psychological Recovery from Work Demands and Employee Mental Health. In *The Routledge Companion to Mental Health at Work* (pp. 267-285). Routledge.
9. Gips, J., Spiegel, A., Norton, A., Gandhi, P., Hardenbergh, D., Gatti, J., Pugh, L., Jones, A., & Sufrin, C. (2020). Health care in the age of mass incarceration: a selective course for medical students in their preclinical years. *MedEdPORTAL*, 16, 11014.
10. Harwood, A. E., Satterfield, K., Helton, W. S., McKnight, P. E., & Shaw, T. H. (2024). The role of state and trait self-control on the sustained attention to response task. *Human factors*, 66(10), 2366-2378.
11. Hasan, M. K. (2024). Digital multitasking and hyperactivity: unveiling the hidden costs to brain health. *Annals of Medicine and Surgery*, 86(11), 6371-6373.
12. Hashmi, S. A., Gundlapalli, R., & Zavar, I. (2025). Mortality in older adults with epilepsy: An understudied entity. *Epilepsia Open*.
13. He, D., Yuan, M., Dang, W., Bai, L., Yang, R., Wang, J., Ma, Y., Liu, B., Liu, S., & Zhang, S. (2023). Long term neuropsychiatric consequences in COVID-19 survivors: Cognitive impairment and inflammatory underpinnings fifteen months after discharge. *Asian Journal of Psychiatry*, 80, 103409.
14. Herlambang, M. B., Cnossen, F., & Taatgen, N. A. (2021). The effects of intrinsic motivation on mental fatigue. *PloS one*, 16(1), e0243754.
15. Hirata, R. P., Thomsen, M. J., Liston, M., Christensen, M. G., & Vestergaard, P. (2024). Clinical features of fallers and non-fallers: finding best-performing combinations of physical performance measurements to discriminate physical impairments between fallers and non-fallers among older adults with and without osteoporosis. *Osteoporosis International*, 35(11), 2007-2016.
16. Joo, H., Diaz-Ramirez, L. G., Chen, C. L., Sun, C. Q., Smith, A. K., Boscardin, W. J., & Whitlock, E. L. (2025). Cognitive Trajectory Before and After Cataract Surgery: A Population-Based Approach. *Journal of the American Geriatrics Society*.
17. Kasprzak, D., Kaczmarek-Majer, K., Rzeźniczak, J., Klamecka-Pohl, K., Ganowicz-Kaatz, T., Słomczyński, M., Budzianowski, J., Pieszek, K., Hiczekiewicz, J., & Tykarski, A.

- (2023). Cognitive impairment in cardiovascular patients after myocardial infarction: Prospective clinical study. *Journal of Clinical Medicine*, 12(15), 4954.
18. Kim, S., Lee, H., Lee, J., Lee, S. W., Kwon, R., Kim, M. S., Koyanagi, A., Smith, L., Fond, G., & Boyer, L. (2024). Short-and long-term neuropsychiatric outcomes in long COVID in South Korea and Japan. *Nature human behaviour*, 8(8), 1530-1544.
 19. Levine, D. A., Gross, A. L., Briceño, E. M., Tilton, N., Kabeto, M. U., Hingtgen, S. M., Giordani, B. J., Sussman, J. B., Hayward, R. A., & Burke, J. F. (2020). Association between blood pressure and later-life cognition among black and white individuals. *JAMA neurology*, 77(7), 810-819.
 20. Li, J., & Carayon, P. (2021). Health Care 4.0: A vision for smart and connected health care. *IIE transactions on healthcare systems engineering*, 11(3), 171-180.
 21. Li, X., Gao, Q., & Liu, Y. (2025). Restoring vigilance in multi-task scenarios: a comparison of meditation, exposure to nature, music, and action video games. *Ergonomics*, 1-23.
 22. Lin, Z., Ye, J., Allore, H., Gill, T. M., & Chen, X. (2024). Early-life circumstances and racial disparities in cognition among older adults in the US. *JAMA internal medicine*, 184(8), 904-914.
 23. Lovett, R., Filec, S., Bonham, M., Benavente, J. Y., O'Connor, R., Russell, A., Zheng, P., Wismer, G., Yoon, E., & Weiner-Light, S. (2023). Long-term impact of the COVID-19 pandemic on self-management of chronic conditions among high-risk adults in the USA: protocol for the C3 observational cohort study. *BMJ open*, 13(10), e077911.
 24. Ma, Y.-H., Chen, H.-S., Liu, C., Feng, Q.-S., Feng, L., Zhang, Y.-R., Hu, H., Dong, Q., Tan, L., & Kan, H.-D. (2023). Association of long-term exposure to ambient air pollution with cognitive decline and Alzheimer's disease-related amyloidosis. *Biological Psychiatry*, 93(9), 780-789.
 25. Mack, M., Stojan, R., Bock, O., & Voelcker-Rehage, C. (2022). Cognitive-motor multitasking in older adults: a randomized controlled study on the effects of individual differences on training success. *BMC geriatrics*, 22(1), 581.
 26. Madore, M. R., Kozel, F. A., Williams, L. M., Green, L. C., George, M. S., Holtzheimer, P. E., Yesavage, J. A., & Philip, N. S. (2022). Prefrontal transcranial magnetic stimulation for depression in US military veterans—A naturalistic cohort study in the veterans health administration. *Journal of Affective Disorders*, 297, 671-678.
 27. McEwen, B. (2024). ALZHEIMER'S DISEASE AND NUTRITIONAL AND HERBAL MEDICINE: A BRIEF LOOK INTO THE INTRICATE METABOLIC NETWORK. *Australasian College of Nutritional & Environmental Medicine Journal*.
 28. McEwen, S., Jarrahi, B., Ventura, J., Subotnik, K., Nguyen, J., Woo, S., & Nuechterlein, K. (2023). A combined exercise and cognitive training intervention induces fronto-cingulate cortical plasticity in first-episode psychosis patients. *Schizophrenia research*, 251, 12-21.
 29. Menon, V., Coppard, M., McEwen, S., Romero, L., Kennedy, E., & Azzopardi, P. (2025). Evaluated Interventions Targeting the Mental Health and Psychosocial Wellbeing of Children and Adolescents: A Scoping Review Focused on Low-and Middle-Income Countries in East Asia and the Pacific. *Asia Pacific Journal of Public Health*, 10105395241313154.
 30. Nelson, A., Aronsson, I., Tillfors, M., Neely, A. S., & Gavelin, H. M. (2025). The experienced route to cognitive health: Cognitive recovery in persons with prior stress-related Exhaustion disorder. *BMC psychiatry*, 25(1), 375.
 31. Nguyen, L. (2021). Carputer with LoRa: Car Location Monitoring System Based on LoRa Network.
 32. O'Hara, L., Neville, C., Marr, C., McAlinden, M., Kee, F., Weir, D., & McGuinness, B. (2024). Investigating the prevalence of cognitive impairment and dementia in the Northern

- Ireland Cohort for the Longitudinal Study of Ageing (NICOLA): the Harmonised Cognitive Assessment Protocol (HCAP) cross-sectional substudy. *BMJ open*, 14(1), e075672.
33. Ophir, G. (2023). *Associations Between Maternal Behaviors and Preschool Children's Stress Reaction During COVID-19-Related Lockdown: Examining the Potential Underlying Role of Child Executive Functioning* Reichman University (Israel)].
 34. Reitmajer, M., Leiter, U., Nanz, L., Amaral, T., Flatz, L., Garbe, C., & Forschner, A. (2024). Long-term survival of stage IV melanoma patients: evaluation on 640 melanoma patients entering stage IV between 2014 and 2017. *Journal of Cancer Research and Clinical Oncology*, 150(1), 15.
 35. Schumann, F., Steinborn, M. B., Kürten, J., Cao, L., Händel, B. F., & Huestegge, L. (2022). Restoration of attention by rest in a multitasking world: Theory, methodology, and empirical evidence. *Frontiers in Psychology*, 13, 867978.
 36. Shaw, B. C., Anders, V. R., Tinkey, R. A., Habean, M. L., Brock, O. D., Frostino, B. J., & Williams, J. L. (2024). Immunity impacts cognitive deficits across neurological disorders. *Journal of Neurochemistry*, 168(10), 3512-3535.
 37. Sonnentag, S., Cheng, B. H., & Parker, S. L. (2022). Recovery from work: Advancing the field toward the future. *Annual Review of Organizational Psychology and Organizational Behavior*, 9(1), 33-60.
 38. Strayer, D. L., Castro, S. C., Turrill, J., & Cooper, J. M. (2022). The persistence of distraction: The hidden costs of intermittent multitasking. *Journal of experimental psychology: applied*, 28(2), 262.
 39. Weller, P. (2022). Effects of Long-Term Memory on Visual Attention and Access to Visual Consciousness.
 40. Westbrook, J., Raban, M. Z., Clive, J., & Walter, S. R. (2025). Interruptions and multitasking in clinical work: a summary of the evidence. In *Reengineering Clinical Workflow in the Digital and AI Era: Toward Safer and More Efficient Care* (pp. 163-177). Springer.
 41. Zhang, S., Zhang, N., Wang, S., Hong, J., Li, F., Guo, H., Lv, Z., Wang, Y., Wang, W., & Wu, W. (2023). Circadian rhythms and sleep quality among undergraduate students in China: the mediating role of health-promoting lifestyle behaviours. *Journal of Affective Disorders*, 333, 225-232.